

Examination Committee Approval

Dedication

Acknowledgements	iv
Abstract	vi
Abstract in Arabic	viii
Table of Contents	x
List of Figures	xiii
List of Tables	xvi
List of Symbols and Terminology	xviii

Chapter I: Introduction

1.1 Pesticides	1
1.1.1 2, 4-Dichlorophenoxyacetic Acid	2
1.1.2 Apoptosis and 2, 4-D.....	5
1.2 Vibrational Spectroscopy	6
1.2.1 Fourier transform infrared (FTIR) spectroscopy	6
1.2.2 Theory of Infrared spectroscopy	7
1.3 Medicinal Plants	10
1.3.1 <i>Lepidium sativum</i>	10
1.4 Aim of the study.....	12

Chapter II: Review of Relevant Literature

2.1 Pesticides toxicity.....	14
2.2 Protective effect of medicinal plants.....	16
2.3 Diagnosis by vibrational spectroscopy.....	17

Chapter III: Materials and Methods

3.1 Chemicals and preparation of <i>Lepidium sativum</i> aqueous extract.....	20
	21
	22
	22

3.2 Animals and studied groups	
3.3 Diet and measurement of body weight, liver weight and weight gain	
3.4 Tissue collection	
3.5 Infrared measurement.....	23
3.6 Statistical analysis.....	24
3.7 Transmission electron microscopy.....	24

Chapter IV: Results

4.1 Brain cerebellum results.....	26	4.1.1
Ratio measurements	31	
4.1.2 Infrared spectral signature	36	
4.1.2.1 The region 3600-3050 cm ⁻¹ : Analysis of Hydroxyl and Hydroperoxyl Bands.....	36	
4.1.2.2 The region 3050-2800 cm ⁻¹ : Symmetric and asymmetric stretching of methyl (CH ₃) and methylene (CH ₂) functional group.....	42	
4.1.2.3 The region 1800-1500 cm ⁻¹ : Amide I and Carbonyl... Analysis of Carbonyl Band	50	
4.1.3 Electron microscopy.....	52	
4.2 Liver Results.....	61	
4.2.1 Infrared spectral signature.....	69	
4.2.1.1 The spectral region 1000cm ⁻¹ – 1200cm ⁻¹	69	
4.2.1.2 The spectral region 1800cm ⁻¹ – 1500cm ⁻¹	71	
4.2.2 Ratio measurements.....	72	
4.2.2.1 The lipid to protein ratio.....	77	

4.2.2.2 Polarity, chain length and packing of membrane lipids ratios; $\text{CH}_2(2925\text{cm}^{-1})/\text{CH}_3(2866\text{cm}^{-1})$, $\text{CH}_2(2925\text{cm}^{-1})/\text{total lipids}$ and $\text{CH}_2(2852\text{cm}^{-1})/\text{CH}_3(2866\text{cm}^{-1})$ ratios.....	77
4.2.2.3 Carbonyl ester/total lipid	78
4.2.2.4 Protein ratio.....	79
4.3 Electron Microscope Results.....	79
4.4 The evaluation of liver weight and body weight	87
Chapter V: Discussion	
5.1 Brain tissue	90
5.2 Liver tissue	99
Chapter VI: Conclusions and Recommendations	
Conclusions and Recommendations	107
List of References	110
Arabic Summary	

LIST OF FIGURES

Figure	Page
1.1 2,4-Dichlorophenoxy acetic acid Chemical Structure.....	3
1.2 (a)Symmetric and asymmetric stretching vibrations, (b) different types of bending vibrations.....	9
1.3 <i>Lepidium sativum</i> seeds.....	13
4.1 Representative overlaid infrared spectra of rat brain tissues obtained from cont., 2, 4-D, water cont and LS treated groups. The entire spectra were normalized and baseline corrected.....	28
4.2a Comparison of the coadded FTIR spectra of cont. and 2, 4-D groups in the mid infrared region (4000-400 cm ⁻¹).	39
4.2b Comparison of the coadded FTIR spectra of water cont. and LS treated groups in the mid infrared region (4000-400 cm ⁻¹).....	30
4.3 The values of protein ratios [I(3303cm ⁻¹)/I(3072 cm ⁻¹)] and [I(1652 cm ⁻¹)/I(2921 cm ⁻¹)] of brain tissues for all groups under investigation	34
4.4 The values of lipid chain packing ratio I(2852 cm ⁻¹)/I(2866 cm ⁻¹) of brain tissues for all groups under investigation.....	34
4.5 The values of lipid polarity ratio [I(2921 cm ⁻¹ /I(2866 cm ⁻¹)]of brain tissues for all groups under investigation	35
4.6 The values of ester (C=O) ratio [I(1735cm ⁻¹)/I(1547cm ⁻¹)] ratio indicative of the weight of formation of carbonyl compounds of brain tissues for all groups under investigation.....	35
4.7a Curve fitting obtained from brain tissues of 2, 4-D and control groups in the IR spectral range (3700-2700cm ⁻¹)	38
4.7b Curve fitting obtained from brain tissues of treated and water control groups in the IR spectral range (3700-2700cm ⁻¹)	39
4.8 Kubelka munk analysis obtained from brain tissues of (a) 3600-3050cm ⁻¹ (b) 3050-2800 cm ⁻¹ (c) 1800-1500cm ⁻¹ IR spectral regions for all treated groups under investigation.....	44

4.9a Deconvoluted and peak resolved IR spectra of the brain tissue over the range (1800-1500cm ⁻¹) for control and 2, 4-D treated groups.....	47
4.9b Deconvoluted and peak resolved IR spectra of the brain tissue over the range (1800-1500cm ⁻¹) for water control and all LS treated groups.....	48
4.10 Transmission electron micrograph of the control cerebellum rat brain	53
4.11 Transmission electron micrograph of the control cerebellum of rat brain showing mitotic division in cytokinesis stage	54
4.12 Transmission electron micrograph of the cerebellum brain of 2, 4-D treated group	55
4.13 Transmission electron micrograph of the cerebellum of rat brain of the curative group.....	56
4.14 Transmission electron micrograph of the cerebellum of rat brain of the protective group	57
4.15 Transmission electron micrograph of the cerebellum of rat brain of the LS only group	58
4.16 Transmission electron micrograph of the cerebellum of rat brain of the water control group	59
4.17a Coadded and overlaid infrared spectra (3050-2800 cm ⁻¹ spectral region) of three rats liver tissues from all treated groups under investigation.....	62
4.17b Coadded and overlaid infrared spectra (1800-1400 cm ⁻¹ spectral region) of three rats liver tissues from all treated groups under investigation.....	62
4.18a Curve fitting obtained from liver tissues of 2, 4-D and control groups in the IR spectral range (3700-2700cm ⁻¹)	64
4.18b Curve fitting obtained from liver tissues of 2, 4-D and control groups in the IR spectral range (3700-2700cm ⁻¹)	65
4.19a Deconvoluted and peak resolved IR spectra of the liver tissue over the range (1800-1500cm ⁻¹) for control and 2, 4-D treated.....	66
4.19b Deconvoluted and peak resolved IR spectra of the liver tissue over the range (1800-1500cm ⁻¹) for water control and all LS treated groups.....	67
4.20 Coadded and overlaid infrared spectra 1400-900 cm ⁻¹ spectral region) of three rats liver tissues from all groups under investigation.....	70
4.21 Area ratios of liver tissue. (a) Carbonyl ester/total lipid ratio at (1735cm ⁻¹ /total lipid) indicating the carbonyl status in the system, (b) Total lipid/total protein at [CH ₂ (2925cm ⁻¹) + CH ₂ (2854cm ⁻¹)] /[(Amide I+ Amide II)], (c)	

vas(CH ₂)lipids/total lipid at (2921cm ⁻¹ /total lipid) indicating the alterations in phospholipids chain length.....	75
4.22 Intensity ratios of liver tissue; (a) Amide A/Amide B at (3301cm ⁻¹ /3072cm ⁻¹) for protein content, (b) v _{as} (CH ₂) / v _s (CH ₃) at (2925cm ⁻¹ /2866cm ⁻¹) for polarity of lipid chains, (c) v _s (CH ₂) / v _s (CH ₃) at (2854cm ⁻¹ /2866cm ⁻¹) for looseness of lipid chains packing.....	76
4.23 Transmission electron micrograph of the control rat liver tissues	81
4.24 Transmission electron micrograph of the rat liver sacrificed 24h after 2,4-D treatment.....	82
4.25 Transmission electron micrograph of the rat liver of curative group	83
4.26 Transmission electron micrograph of the rat liver of protective group	84
4.27 Transmission electron micrograph of the rat liver of LS only group	85
4.28 Transmission electron micrograph of the rat liver of water control group.....	86